Table 29.6

5-Year Relative Survival (Percent), 2006-2012^a

By International Classification of Childhood Cancer(ICCC)^b Selected Group and Subgroup^c and Sex and Age Excluding benign brain and myelodysplastic syndromes

	Ages 0-19 by Sex			Both Sexes by Age					
ICCC Group & Subgroup	Total	Male	<u>Female</u>	<1	1-4	5-9	10-14	15-19	0-14
All ICCC Groups Combined	83.3	82.1	84.6	78.7	84.5	83.2	82.3	83.9	83.0
I Leukemia	83.1	82.8	83.4	62.0	90.0	89.3	77.8	71.3	85.6
I(a) Lymphoid leukemia	88.1	87.5	88.8	60.8	93.7	92.6	82.2	74.7	90.2
I(b) Acute myeloid leukemia	62.9	62.4	63.3	60.0	66.5	69.1	61.2	59.7	64.2
II Lymphomas and reticuloendothelial neo.	93.1	92.9	93.2	79.9 ^d	92.6	94.1	93.8	92.7	93.4
II(a) Hodgkin lymphoma	96.9	96.7	97.0	-	94.5 ^d	97.7	97.9	96.4	97.7
II(b,c,e) Non-Hodgkin lymphoma	88.8	89.3	87.7	-	91.3	92.2	89.7	86.0	90.7
III CNS & misc intracranial & intraspinal neo. III(a) Ependymomas and choroid plexus tumor III(b) Astrocytoma III(c) Intracranial and intraspinal embryonal tumors III(d) Other gliomas	73.8 77.9 82.6 62.8 59.9	72.9 77.3 81.1 62.3 63.4	74.8 78.6 84.1 63.5 56.2	60.3 65.3 ^e 81.8 29.1	72.8 74.7 90.6 55.6 49.2	70.6 77.5 ^d 80.5 70.6 44.7	78.6 82.1 ^d 81.3 76.8 70.4	79.1 88.1 77.2 73.2 83.3	72.6 75.9 83.9 61.5 54.1
IV Neuroblastoma and other peripheral nervous cell tumors $\ensuremath{\text{IV}}(a)$ Neuroblastoma and ganglioneuroblastoma	79.5 79.5	78.0 78.2	81.2 81.0	93.8 93.8	73.8 73.7	70.2 69.8	72.3 ^d 72.6 ^d	74.2 ^d	79.7 79.6
V Retinoblastoma	95.3	94.2	96.5	97.0	93.5	-	-	-	95.3
VI Renal tumors	89.2	87.2	91.1	94.0	91.0	90.1	76.3 ^d	68.1 ^d	90.6
VI(a) Nephroblastoma and other nonepithelial renal tumor	91.1	89.3	92.8	94.0	91.2	91.7	82.1 ^e		91.5
VII Hepatic tumors VII(a) Hepatoblastoma	72.4 83.9	71.3 82.0	74.6 87.2	86.3 85.9	83.1 83.1	67.5 ^d	35.4 ^e	47.4 ^d	77.1 84.2
VIII Malignant bone tumors VIII(a) Osteosarcoma VIII(c) Ewing tumor and related sarcomas of bone	70.7	69.1	73.0	-	80.2 ^d	77.6	72.2	65.7	74.2
	67.0	65.1	69.4	-	-	72.4	68.8	63.4	69.5
	71.7	68.8	76.9	-	94.6	83.0	72.9	59.2	78.7
IX Soft tissue and other extraosseous sarcomas IX(a) Rhabdomyosarcoma	72.3	70.5	74.5	64.5	75.8	78.8	72.6	69.1	74.0
	65.8	66.5	64.9	53.0 ^d	72.7	78.7	59.2	48.9	69.6
X Germ cell & trophoblastic tumors & neo. of gonads $X(a)$ Intracranial and intraspinal germ-cell tumor $X(c)$ Malignant gonadal germ cell tumor	92.4	92.8	91.6	87.7	93.3	98.3	94.0	91.9	93.3
	90.0	91.1	85.6 ^d	-	-	98.4	91.1	89.2	90.4
	96.7	96.1	97.8	100.0	100.0	98.0	99.4	95.7	99.4
XI Other malignant epithelial neo. and melanomas $^{\rm f}$ XI(b) Thyroid carcinoma XI(d) Malignant melanoma	92.2	86.6	94.9	-	96.3	93.2	93.0	91.8	92.9
	99.7	98.9	99.8	-	-	98.1	100.0	99.7	99.7
	93.9	91.1	96.0	-	96.3	96.0	94.2	94.0	93.7

a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Based on follow-up of patients into 2012.

International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005: Vol 103, No. 7, pg 1457-1467.

Classifications are shown for invasive cases only. For ICCC groups and subgroups that include in situ behavior, only invasive cases are shown except as noted.

The standard error is between 5 and 10 percentage points.

The standard error is greater than 10 percentage points.

Includes in situ urinary bladder tumors.

⁻ Statistic could not be calculated due to fewer than 25 cases during the time period. Neoplasms are abbreviated as neo.